



# An Alternative Approach to Rodent Control

 **BASF**  
We create chemistry

## Anticoagulants and the rise of the resistant rodent.

Throughout the years, there have been many innovations focused on making anticoagulant rodent baits work faster and more efficiently. Anticoagulant rodent baits cause disruption of clotting activity three to four days after ingestion; this leads to internal bleeding, a coma, and the subsequent death of the rodent. Anticoagulants have been used for many years due to their effectiveness. They also have the ability to circumvent the taste-aversion phenomenon (due to the delayed action) present in many rodents.

After heavy use of first-generation anticoagulants (warfarin, coumatetralyl, chlorophacinone, diphacinone), resistance cases were described for the first time in Scotland in 1958 (Lund, 1972). The anticoagulants stopped working in practical conditions when bait was applied correctly. A genetic phenomenon linked to mutations of the *VKORC1* gene in rodents allowed them to remain unaffected by these anticoagulants. The rodents were no longer susceptible to these compounds (Greaves, 1994). The genetic basis of resistance was described years later (Rost et al., 2004). The development of second-generation anticoagulants (bromadiolone, difenacoum, flocoumafen, difethialone, brodifacoum) made it possible to overcome this resistance. However, these second-generation anticoagulants all act according to the same mechanism and many of these molecules can lead to genetic resistance in rodents. Resistance to bromadiolone and difenacoum has already appeared in brown rats, black rats and mice. The good news is that resistance to second-generation molecules is far more geographically limited than resistance to first-generation anticoagulants.


## Recent trends

The use of second-generation molecules, especially the newest molecules of the second generation (anticoagulants, brodifacoum, flocoumafen, and difethialone), has increased and they still work well. Nevertheless, it is crucial to continuously introduce molecules with different modes of action to prevent new resistances from developing in these mammals. Rodent control professionals need alternatives to anticoagulants to complement the current control solutions available for a long term solution.



**above:**

*If rodents are seen during the day the colony may be so well established that they no longer feel threatened foraging in the daylight.*



**above:**

*Chemical structure of cholecalciferol*





## Possible solutions

There are currently four alternatives to the anticoagulant resistance problem that are considered credible. One is alphachloralose, which acts on rodents' nervous systems and becomes toxic between seven and 12 hours after the bait is consumed. It is extremely effective on mice, but due to poor efficacy against rats alphachloralose is not registered for use against this pest. It is also restricted to indoor use in Europe. Another is cholecalciferol (Vitamin D3). It can be used as a rodenticide and is effective on both rats and mice. This molecule acts on calcium homeostasis, and initially causes the animal to stop consuming the bait quickly ("stop-feeding" effect). The toxic symptoms and death occur two to three days after consuming the bait, which is enough time to prevent the rat from becoming distrustful. And cholecalciferol is widely accepted by rodents because it is not easy to detect and generates no resistance phenomenon known to date. Zinc phosphide is an acute toxicant and has been widely used in managing mouse populations. It causes cardiac and pulmonary dysfunction after only 15 minutes, but is ineffective with rats who are suspicious of rapid poisons. Also, zinc phosphide is not an approved PT 14 biocide in Europe (only aluminum phosphide). Zinc phosphide is a plant protection active for use against voles (*Microtus agrestis* and *Clethrionomys glareolus*). Finally, bromethalin is a neurotoxic agent that has never been used in Europe but is widely used in the United States against rats and mice. Its action is time-delayed like cholecalciferol, but the symptoms suggest significant suffering in animals.

These alternative active ingredients, in particular cholecalciferol (in rats and mice) and alphachloralose (in mice), slow the development of resistance to second-generation anticoagulants and create a unique set of tools to manage rodent populations.

## Summary

Humans have always struggled to keep rodents out of our homes and businesses. Sanitation is the primary reason for rodent control. In fact, brown rats, black rats and mice are capable of transmitting hundreds of infectious diseases to humans, including leptospirosis, hantaviruses, and numerous pathogenic bacteria (Lasseur et al., 2007). With the risk of resistance to second-generation anticoagulants, it is critical that we develop alternatives now to increase the modes of action available in our rodent control toolbox.

### About BASF's Crop Protection division

*With a rapidly growing population, the world is increasingly dependent on our ability to develop and maintain sustainable agriculture and healthy environments. BASF's Crop Protection division works with farmers, agricultural professionals, pest management experts and others to help make this possible. With their cooperation, BASF is able to sustain an active R&D pipeline, an innovative portfolio of products and services, and teams of experts in the lab and in the field to support customers in making their businesses succeed. In 2016, BASF's Crop Protection division generated sales of more than €5.6 billion. For more information, please visit us at [www.agriculture.basf.com](http://www.agriculture.basf.com) or on any of our social media channels.*

BASF 26 Davis Drive, Research Triangle Park, NC 27709 USA, [rodentcontrol@basf.com](mailto:rodentcontrol@basf.com)  
© 2017 BASF Corporation. All Rights Reserved. GL-XXXX August 2017



### above:

*Romain Lasseur, Toxicologist (PhD, MBA), CEO & Founder of Izipest. Rodent and insect expert with more than 15 years of experience in pest management. Romain has overseen many international pest management projects and is deeply involved in designed pest management innovation and solutions. He now works to educate pest control users and industries.*

### Sources

**RRAC: Rodent Resistance Action Committee**  
<http://rrac.info>  
<http://rrac.info/content/uploads/RRAC%20Guidelines-Rodenticide-Resistance-Sept2015.pdf> [http://rrac.info/content/uploads/RRAC\\_Guidelines-Rodenticide-Resistance-Sept2015.pdf](http://rrac.info/content/uploads/RRAC_Guidelines-Rodenticide-Resistance-Sept2015.pdf)

**Lund, M. (1972).**  
Rodent resistance to the anticoagulant rodenticides with particular reference to Denmark.  
Bull. WHO, 47, 611-618.

**Greaves, J. (1994)**  
Resistance to anticoagulant rodenticides.  
In: Buckle, A.P., Smith, R.H. (eds). Rodent pests and their control, Cabi Publishing, Cambridge, 194-217

**Lasseur, R., Longin-Sauvageon, C., Videmann, B., Billeret, M., Berry, P., Benoit, E. (2005).**  
Warfarin resistance in a French strain of rats.  
J. biochem. mol. Toxicol., 19, 379-385

**Rost, S., Fregin, A., Ivaskевичius, V., Conzelmann, E., Hörtnagel, K., Pelz, H.-J., Lappegard, K., Seifried, E., Scharrer, I., Tuddenham, E., Müller, C., Strom, T., Oldenburg, J. (2004).**  
Mutations in *VKORC1* cause warfarin resistance and multiple coagulation factor deficiency type 2.  
Nature, 427, 537-541.

  
We create chemistry